

The Board						
Dr F. Corrad President	Pr R. Cohen Scientific Director	Dr M. Boucherat Treasurer	Dr C. Batard Ass. Treasurer	Dr A. Elbez Secretary		
Dr A-S. Michot	Dr O. Romain	Dr C. Schlemmer	Dr E. Seror	Dr C. Turberg-Romain	Dr A. Werner	Dr A. Wollner

The Team	
Dr C. Levy Medical Director	I. Destel, M. Servera Project Manager
I. Ramay Executive Assistant	P. Daumerie Clinical Research Associate
S. Béchet Biometric Manager	A. Prieur, J. Chalte Clinical Research Assistant


The Partner Laboratories	
Microbiology department Hôpital Robert Debré (Paris)	Biologie department CHI de Créteil
Pr S. Bonacorsi, PU-PH, Department Head and Associate Laboratory Director of CNR E. Coli et Shigella	Dr E. Varon, PH, Head of Department and Director of the National Reference Center for Pneumococcus
E. Sobral Laboratory technician	M. Lorin Laboratory technician

The Research

ACTIV has the **experience** as well as structural, staff and methodological means to perform clinical studies, **from protocol conception to national and international publication.**

These studies are initiated by ACTIV, pediatricians in the field, university and official agencies, and the pharmaceutical industry. They cover research into **infectious diseases** (observatory, epidemiology, diagnostic, therapeutic methods, vaccinology) and aim to contribute to **“evidence-based medicine”**.

Although most of the studies are performed in an **out-of-hospital** environment, some take place in a **hospital environment** at a regional or national level.



WHO ARE WE ?

1988
ACTIV
Develop research in ambulatory paediatric
Anti-infectious and vaccines studies

2001
GPIP
Pediatric Infectious Pathology Group of the French Pediatric Society

2003
INFOVAC FRANCE
Information in vaccinology

2012
CRC Créteil (2012)
GRC GEMINI
Créteil University

WHO ARE ACTIV'S PARTNERS?

ACTIV WORKS WITH

1000
AMBULATORY PAEDIATRICIANS /AFPA

250
SERVICES OF HOSPITAL PAEDIATRICIANS AND MICROBIOLOGISTS

BY WHOM ARE THE SAMPLES ANALYZED IN THE STUDIES?


NATIONAL REFERENCE CENTRES

HOW ARE ACTIV STUDIES FUNDED ?

GRANTS
INDUSTRY (ATB AND VACCINES)
PHRC
OFFICIAL AGENCIES

STUDIES WITHOUT GRANTS
ACTIV

ACTIV
31 rue Le Corbusier
94 000 Créteil
+33 (0)1.48.85.04.04
activ@activ-france.fr
<https://www.activ-france.com/en/>



The Expertise




The Network

The network of investigators involves a **core of 100 pediatricians**, however, for some studies (for example on flu tests), **this number can reach 1,000**. The hospital network consists of **250 pediatric and 170 microbiology services**.

The Publications

ACTIV publishes **annually** between 20 and 30 articles in peer-reviewed journals. You can find all our publications on [our website](#), in the section " Publications ".



Evolution of ACTIV publications since it beginning in 1988

Pediatric Ambulatory and Hospital Networks for Surveillance and Clinical Epidemiology of Community-Acquired Infections

Corinne Levy, MD^{1,2,3,4,5}, François Vie le Sage, MD^{4,5}, Emmanuelle Varon, MD⁶, Martin Chalumeau, MD^{4,7,8}, Emmanuel Grimprel, MD^{4,9}, and Robert Cohen, MD^{1,2,3,4,5,10}

Care of infectious diseases (prevention, diagnosis, and treatment) represents a large part of the activity of pediatric practices as well as primary care, emergency departments, and hospitals. The bacterial and viral species involved in pediatric community-acquired infections (CAIs) can induce severe, moderate, or mild diseases, which require care by hospitalization, emergency departments, or first-line clinicians. For example, pneumococcus manifests as a spectrum of diseases ranging from severe invasive diseases, such as meningitis, bacteremic pneumonia, and bacteremia, to less severe but more frequent diseases, such as acute otitis media (AOM), sinusitis, and pneumonia (Figure; available at www.jpeds.com).^{1,2} Therefore, outpatient use of antibiotics also involves antibiotic resistance for CAIs; hence, infection management becomes even more complex, often leading to hospitalizations. Moreover, some vaccines, for which the main objective is reducing invasive and severe diseases, have an additional impact on less severe diseases and also change the carriage.³ Trying to study the bacterial and clinical epidemiology of organisms such as *Pneumococcus*, Group A *Streptococcus*, or *Escherichia coli*, involved in a large spectrum of CAIs, requires a focus on both the ambulatory setting and on the hospital.

We review the efforts to build a French pediatric research network that focused on CAIs.

Before the 1990s, in France, first-line clinicians and notably ambulatory pediatricians were not involved in clinical research, which was performed mainly by universities and hospitals. Because of a lack of surveillance systems in ambulatory settings, we created a nonprofit ambulatory-pediatric research network, Pediatric Clinical and Therapeutic Association of Val de Marne (ACTIV). This regional network (Paris area) was extended at the national level via a strong collaboration with the Association of French Ambulatory Pediatricians. A link with a preexisting hospital network, the Pathology Pediatric Infectious Disease Group of the French Pediatrics Society, was established several years later. The network has complied with the high-quality standards required by good clinical practice for industrial trials by European and North American drug regulators and the “feasibility in real life,” taking into account medical practice with ambulatory care and

hospital constraints. With ACTIV, we have designed study protocols aimed at simplifying the procedures, while maintaining a high standard of quality. This approach facilitated the publication of scientific data that were widely used by the European and American medical markets.

From Antibiotic Treatment to Prevention with Vaccines

To compare the efficacy of different therapeutic options (type of drug, dosage, duration of antibiotic treatments, etc), in AOM we have performed several clinical trials using rigorous criteria to standardize the AOM diagnosis along with otoscopy training sessions.⁴⁻⁶ Our research group evaluated the impact of antibiotics on the composition and antibiotic resistance in nasopharyngeal flora that involved a network of almost 100 primary care pediatricians.^{5,6} The bacteriological samples were centralized in the National Reference Center for Pneumococci. The advantage of the expertise acquired in this area (several thousand nasopharyngeal samples collected in 10 years) allowed us to perform our pivotal study on nasopharyngeal carriage after the 7-valent pneumococcal conjugate vaccine (PCV) implementation in France in 2001.⁷ This study was performed as a postlicensing commitment requested by the European Medicines Agency to determine the impact of the 7-valent pneumococcal conjugate vaccine. Although other studies reported similar results for the PCV impact, none were comparable with those we have conducted since 2001 in terms of design, duration (>17 years), and number of patients enrolled (>15 000).⁷

ACTIV	Pediatric Clinical and Therapeutic Association of Val de Marne
AOM	Acute otitis media
CAI	Community-acquired infection
PCV	Pneumococcal Conjugate Vaccine

From the ¹University Paris Est, IMRB- GRC GEMINI, Créteil, France; ²ACTIV, Pediatric Clinical and Therapeutic Association of the Val de Marne, Saint-Maur des Fossés; ³Clinical Research Center (CRC), Centre Hospitalier Intercommunal de Créteil; ⁴GPIP, Pediatric Infectious Disease Group; ⁵AFFPA, French Association of Ambulatory Pediatricians, Saint-Germain-en-Laye; ⁶National Reference Center for Pneumococci, Microbiology Laboratory, Assistance Publique-Hôpitaux de Paris, Hospital Georges-Pompidou; ⁷Obstetrical, Perinatal and Pediatric Epidemiology Research Team (Epopé), Center for Epidemiology and Statistics Sorbonne Paris Cité (CRESS-INSERM U1153), Paris Descartes University; ⁸Department of General Pediatrics and Pediatric Infectious Diseases, Necker hospital for Sick Children, Assistance Publique-Hôpitaux de Paris, Paris Descartes University; ⁹Department of General Pediatrics, Hôpital Trousseau, AP-HP, University Pierre et Marie Curie, Paris; and ¹⁰Unité Court Séjour, Petits Nourrissons, Neonatology Department, Centre Hospitalier Intercommunal de Créteil, France

The institutions of C.L. and R.C. (ACTIV) received research grant support from Pfizer, GlaxoSmithKline, and Sanofi Pasteur MSD outside of the submitted work. The authors declare no conflicts of interest.

0022-3476/\$ - see front matter.
<https://doi.org/10.1016/j.jpeds.2017.11.050>

Keys to Success

Several factors could explain the success of this network lasting more than 30 years (Table; available at www.jpeds.com). The important outcome of the studies performed by the research network has been the implementation of clinical conclusions and recommendations by pediatricians in their practice. This includes the duration of antibiotic therapy for children with AOM or group A *Streptococcus* pharyngitis as well as the use of biomarkers or rapid diagnostic tests to improve diagnostic performance and use of appropriate antibiotics for different infectious diseases.⁸⁻¹¹ Finally, the best proof of success was that pediatricians' behavior was correlated with the research they conducted. Particularly, our group, which promptly implemented the proposed recommendations, showed less prescription of antibiotics and greater vaccination coverage for their patients than other physicians who applied the recommendations later.¹²

We have recently moved to a new method that allows us to even more easily perform studies by directly obtaining data from pediatricians' computers. With 100 pediatricians using the same software (Infansoft, CompuGroup Medical, Koblenz, Germany), we have automated data capture from electronic medical records for children in ambulatory settings, the Panel in Ambulatory Research Infectiology. The participants benefit in real time from the epidemiology of several infectious diseases on a dedicated Website. They can also improve their diagnosis with e-learning sessions specifically dedicated to each pathology.

Link with the Hospital Network

Linking ambulatory and hospital networks is important. For instance, to determine the multifaceted impact of pneumococcal vaccine implementation on invasive pneumococcal infections, we linked the ambulatory with the hospital-based surveillance systems by reconciling separate databases, and we created a national hospital network for bacterial meningitis with the Pathology Pediatric Infectious Disease Group of the French Pediatrics Society.^{13,14} More than 230 pediatricians and 168 microbiologists nationally were involved in this study. Microbiologists agreed to voluntarily send each bacterial species isolated from meningitis samples to the corresponding national reference center, which performed serotyping and genotyping, as well as extensive standardized susceptibility testing. The strong participation and motivation of hospital pediatricians and microbiologists were related to the lack of a prior surveillance system that included clinical, therapeutic, and microbiological data for invasive diseases in France. Moreover, the originality and the relevance of our system lies in the fact that we used our research platform that was initially created for outpatient infectious diseases. We identified a team of pediatricians and microbiologists in each participating center and organized a close collaboration with the experts of the national reference centers for the different bacterial species. The

research has improved the quality of care, and the network has been regularly asked to perform ancillary studies and specific analyses for each bacteria involved. The bacterial meningitis study surveillance, with more than 6500 cases enrolled since 2001, allowed for the publication of several articles, also involving the contribution of young pediatricians.¹⁵ Moreover, our ambulatory and hospital network initially built for pneumococcal infections was extended to study other diseases and pathogens, such as the increasing incidence of CAIs owing to extended-spectrum β -lactamase-producing *E. coli*. These studies provided useful public health data and recommendations.

French Vaccine Network

In 2003, following the Infovac-Swiss model, we developed InfoVac-France, a website designed by Clair-Anne Siegrist of the University of Geneva, providing physicians with a direct source of information on vaccinations.¹⁶⁻¹⁸ In the context of vaccine hesitancy in the world and particularly in France, here again this network helps provide optimal vaccine support and represents a good opportunity for the release of validated scientific information on vaccines.

Conclusion

Rather than providing guidance for better surveillance of pediatric infectious diseases, herein we present an overview of our surveillance system for diseases and highlight why, in the French context, it was successful beyond our hopes. Owing to different clinical practices and healthcare systems, epidemiology, and vaccination programs worldwide, results from studies performed in other countries cannot always be extracted or transposed to one's own country. This factor has led to each country performing its own clinical research. The strength of our surveillance systems lies in the multiplicity of funding (governmental, university, and pharmaceutical industry). Our challenges are to rigorously maintain several well-established surveillance systems with resources specifically allocated and increase the involvement of young researcher-pediatricians. Our networks are not always representative of all French pediatricians, because our researcher-pediatricians are well-informed and particularly well-trained to follow the latest recommendations. However, we believe in this model, which can easily evolve with themes of research prioritized according to epidemiologic changes. Without competing with the different pre-existing research groups in our country, we have federated and involve all the volunteers in our networks. Hence, this "ambulatory/hospital" research network, which is unique in Europe, contributes greatly to answering public health questions, particularly for vaccination strategies and antibiotic resistance.¹⁵ ■

References available at www.jpeds.com

References

1. Bogaert D, De Groot R, Hermans PW. Streptococcus pneumoniae colonisation: the key to pneumococcal disease. Lancet Infect Dis 2004;4:144-54.
2. Simell B, Auranen K, Kayhty H, Goldblatt D, Dagan R, O'Brien KL, et al. The fundamental link between pneumococcal carriage and disease. Expert Rev Vaccines 2012;11:841-55.
3. Dagan R. The diversity of pneumococcal conjugate vaccine impact observed through their implementation. Hum Vaccin Immunother 2016;12:266-7.
4. Cohen R, Levy C, Boucherat M, Langue J, de La Rocque F. A multicenter, randomized, double-blind trial of 5 versus 10 days of antibiotic therapy for acute otitis media in young children. J Pediatr 1998;133:634-9.
5. Varon E, Levy C, De La Rocque F, Boucherat M, Deforche D, Podglajen J, et al. Impact of antimicrobial therapy on nasopharyngeal carriage of Streptococcus pneumoniae, Haemophilus influenzae, and Branhamella catarrhalis in children with respiratory tract infections. Clin Infect Dis 2000;31:477-81.
6. Cohen R, Bingen E, Varon E, de La Rocque F, Brahim N, Levy C, et al. Change in nasopharyngeal carriage of Streptococcus pneumoniae resulting from antibiotic therapy for acute otitis media in children. Pediatr Infect Dis J 1997;16:555-60.
7. Cohen R, Varon E, Doit C, Schlemmer C, Romain O, Thollot F, et al. A 13-year survey of pneumococcal nasopharyngeal carriage in children with acute otitis media following PCV7 and PCV13 implementation. Vaccine 2015;33:5118-26.
8. Cohen R, Reinert P, De La Rocque F, Levy C, Boucherat M, Robert M, et al. Comparison of two dosages of azithromycin for three days versus penicillin V for ten days in acute group A streptococcal tonsillopharyngitis. Pediatr Infect Dis J 2002;21:297-303.
9. Cohen JF, Cohen R, Bidet P, Levy C, Deberdt P, d'Humières C, et al. Rapid-antigen detection tests for group a streptococcal pharyngitis: revisiting false-positive results using polymerase chain reaction testing. J Pediatr 2013;162:1282-4, 4 e1.
10. Cohen R, Levy C, Chalumeau M. Shortened antimicrobial treatment for acute otitis media. N Engl J Med 2017;376:e24.
11. Levy C, Biscardi S, Dommergues MA, Dubos F, Hees L, Levieux K, et al. Impact of PCV13 on community-acquired pneumonia by C-reactive protein and procalcitonin levels in children. Vaccine 2017;35:5058-64.
12. Levy C, Pereira M, Guedj R, Abt-Nord C, Gelbert NB, Cohen R, et al. Impact of 2011 French guidelines on antibiotic prescription for acute otitis media in infants. Med Mal Infect 2014;44:102-6.
13. Cohen R, Biscardi S, Levy C. The multifaceted impact of pneumococcal conjugate vaccine implementation in children in France between 2001 to 2014. Hum Vaccin Immunother 2016;12:277-84.
14. Bingen E, Levy C, de la Rocque F, Boucherat M, Varon E, Alonso JM, et al. Bacterial meningitis in children: a French prospective study. Clin Infect Dis 2005;41:1059-63.
15. Association Clinique et Thérapeutique Infantile du Val de Marne (ACTIV). http://activ-france.com/Publis_en.php. Accessed December 18, 2017.
16. InfoVac Suisse. La ligne directe d'information sur la vaccination! <https://www.infovac.ch/fr/>. Accessed December 18, 2017.
17. InfoVac-France. La ligne directe d'information et de consultation sur les vaccinations. <https://www.infovac.fr/>. Accessed December 18, 2017.
18. Cohen R, Siegrist CA. Vaccines' adverse events: questions and answers brought by InfoVac. Arch Pediatr 2006;13:650-2.




Figure. Spectrum of Streptococcus pneumoniae disease.

Table. Factors determining the successful outcome of pediatric ambulatory and hospital networks for surveillance and clinical epidemiology of CAIs

Addressing areas of typical concern for primary care physicians (ie, improvement in diagnosis and/or management of pediatric infectious diseases).
Developing projects aimed at providing answers to common clinical question related to professional practice and promoting pediatric best practice for the benefit of children (ie, studies involving respiratory tract infections, owing to their frequency and easiness in obtaining oropharyngeal or nasopharyngeal samples).
Placing investigators and their training programs at the cornerstone of the research system (investigators easily recognize that contributing to the projects proposed provides a real benefit to their daily practice and professional continuing education).
Building and establishing preliminary relationships between parents and their family pediatricians, which largely favor parental adherence to protocols and facilitate the collection of parental written informed consent to the studies (very few patients are lost to follow-up).